

Causes, Effects, and Solutions to the Problem of Malnutrition in Older Adults Who Remain Independent

Xuan Tu Nguyen, Student, Gerontology, Hanoi Medical University, Hanoi, Vietnam

Tung Tran, Lecture, Gerontology, Hanoi Medical University, Hanoi, Vietnam

Abstract

Up to 10% of older individuals who live alone experience malnutrition, which raises the risk of morbidity and mortality. This is a severe public health concern. This work investigates the two pathogenic processes associated with anorexia, namely, inflammation-induced tissue catabolism and food deprivation. Reduced appetite, barriers to food access, and socioeconomic issues contribute to nutrient scarcity, impairing physical and cognitive abilities. Chronic inflammation and anorexia are involved in inflammation-induced tissue catabolism, which exacerbates malnutrition and adverse health effects. Malnutrition can reduce bone density, muscle atrophy, immune system weakness, and mental health problems. A multimodal strategy is needed to address malnutrition, including physical activity programmes, medical and psychological care, community support programmes, and nutritional education and counselling. These tactics seek to lessen the load on people and healthcare systems while enhancing senior citizens' health and quality of life. This thorough review emphasizes the frequency, causes, outcomes, and countermeasures to malnutrition's harmful impacts on older persons living independently.

Keywords: malnutrition, older adults, inflammation-induced tissue catabolism, food deprivation, reduced appetite, socioeconomic challenges, chronic inflammation

Introduction

About 10% of older persons who live independently are affected by malnutrition, which is a common problem. Significant health consequences are linked to the illness, such as elevated morbidity and mortality rates. Developing successful therapies requires understanding the underlying mechanisms, such as inflammation-induced tissue catabolism and nutritional deprivation. This study examines how these pathways affect older individuals' health and well-being.

The frequency of malnourishment in the elderly

Malnutrition affects a sizable portion of autonomous older adults. Studies show that between 5 and 10% of these elderly individuals are underweight, and up to 60% of them are at risk of malnutrition. As people age, malnutrition becomes more prevalent, with those over 75 years old being particularly vulnerable.

Pathologic Pathways of Malnutrition

Nutrient Deprivation

Nutrient deficiency results from older persons not consuming enough of certain vital nutrients. This may result from several things, including decreased appetite, barriers to food access, or socioeconomic difficulties. Lack of certain nutrients impairs mental and physical abilities, lowering the overall quality of life.

1. **Reduced Appetite:** Age-related changes often result in a reduced appetite. Alterations in taste and smell can make food less appealing. Dental problems can make chewing difficult, and gastrointestinal issues can lead to discomfort after eating, all contributing to a lower food intake.
2. **Socioeconomic Challenges:** Many older adults face limited income, which can restrict their ability to purchase nutritious foods. Lack of transportation can make it challenging to access grocery stores, and social isolation can reduce the motivation to prepare and consume meals.

Inflammation-Induced Tissue Catabolism and Anorexia

The disintegrating of bodily tissues due to persistent inflammatory reactions is known as inflammation-induced tissue catabolism. Anorexia, a disorder that is widespread in older persons and is characterized by appetite loss, is frequently combined with this process. Combined, these variables worsen malnutrition, which worsens health and makes people more vulnerable to infections and other diseases.

1. **Prolonged Irritation:** "Inflammaging" is the term used to describe the low-grade, persistent inflammation that is linked to ageing. This disease makes it easier for muscles and other tissues to break down, which results in muscle atrophy and a decline in general physical health.
2. **Anorexia of Aging:** Anorexia of ageing is a complex condition influenced by hormonal changes, reduced physical activity, and psychological factors. Hormonal changes can alter hunger signals, reduced physical activity can decrease appetite, and psychological factors such as depression or anxiety can further reduce the desire to eat.

Consequences of Malnutrition in Older Adults

Older people who are malnourished experience a wide range of health problems that negatively affect their general health and raise the possibility of numerous complications. The main repercussions include reduced bone density, diminished immune system performance, muscle atrophy, and adverse mental health effects. These factors increase the chance of fractures and falls, and healing from diseases and procedures takes longer.

Weakened Immune Function

Malnutrition impairs the body's immune response, making older adults more susceptible to infections and delaying their recovery from illnesses. Here is how:

- **Nutrient Deficiencies:** Vital components like manganese and zinc, as well as vitamins A, C, E, and B6, are essential for sustaining effective immunity. Inadequate intake of these nutrients may

impair the body's capacity to generate and stimulate immune cells, reducing the strength of its immune system as a whole.

- **Protein-Energy Malnutrition:** Adequate protein intake is vital for producing antibodies and other immune-related molecules. The body's ability to fight off infections diminishes when insufficient protein intake increases disease vulnerability.
- **Impact on Recovery:** A weakened immune system increases susceptibility to infections and prolongs recovery from illnesses and surgeries. The body's ability to repair tissues and recover from trauma is compromised, leading to more extended hospital stays and increased healthcare costs.

Muscle Wasting and Sarcopenia

Sarcopenia, or the wasting of muscles, is one of the significant effects of malnutrition on the elderly population. The gradual decrease of muscle mass and associated power may significantly impact mobility and independence.

- **Inadequate Protein Intake:** Protein is essential for maintaining muscle mass. The ability to synthesize muscle protein decreases in older adults, making adequate protein intake even more critical. When insufficient protein intake, muscle protein breakdown exceeds synthesis, leading to muscle wasting.
- **Chronic Inflammation:** Chronic low-grade inflammation joint in older adults accelerates muscle protein breakdown. This condition, known as "inflammaging," exacerbates muscle loss and contributes to sarcopenia.
- **Reduced Physical Activity:** Malnutrition often leads to fatigue and a lack of energy, reducing physical activity levels. Physical inactivity further accelerates muscle wasting, as muscles require regular use to maintain their mass and strength.
- **Consequences of Muscle Wasting:** Sarcopenia is associated with decreased strength and mobility, increasing the risk of falls and fractures. It also leads to a loss of independence, as everyday activities such as walking, climbing stairs, and carrying objects become more challenging.

Bone Health

However, malnutrition significantly affects bone health, particularly vitamin D and calcium deficiency, which raises the risk of fractures.

- **Calcium and Vitamin D Deficiencies:** The main building block of bone is calcium-rich soil, and the digestion of calcium requires vitamin D. Lack of these nutrients results in a reduction in bone density, which makes bones brittle and more prone to breaking.
- **Osteoporosis:** Prolonged malnutrition can lead to osteoporosis, a condition characterized by porous and brittle bones. Older adults with osteoporosis are at a significantly higher risk of fractures, even from minor falls or injuries.
- **Increased Fracture Risk:** Serious complications, such as long-term discomfort, impairment, and an increased risk of death, can arise from fractures in the elderly. Particularly concerning are the high rates of death and morbidity linked to broken hips, which might be attributed to consequences like infections and the need for extended immobilization.

Mental Health

The mental health of an individual can be negatively impacted by malnutrition, which can contribute to illnesses such as anxiety and cognitive deterioration.

- **Nutrient Deficiencies:** Essential nutrients, such as oily fish and B Vitamins, along with antioxidant compounds, are necessary for maintaining healthy brain function. Insufficiencies in these nutrients can harm cognitive function and can also raise the chance of developing mental health disorders.
- **Stress of Chronic Conditions:** Living with chronic malnutrition can be psychologically stressful, contributing to feelings of hopelessness and depression. The physical limitations imposed by malnutrition-related conditions can also lead to social isolation, further exacerbating mental health issues.
- **Cognitive Impairment:** Cognitive functions, including memory, concentration, and the decision-making process, could deteriorate due to malnutrition. A decline in quality of life can occur when people with cognitive impairment cannot carry out even the most basic daily tasks.

Interventions to Combat Malnutrition

Addressing malnutrition in independent-living older adults requires a multifaceted approach:

1. **Nutritional Education and Counseling:** Educating older adults on the importance of balanced diets and guidance on meal planning can help improve nutrient intake.
2. **Local Support Programs:** Initiatives such as meal delivery services and communal dining programs can enhance access to nutritious foods.
3. **Medical and Psychological Support:** Regular health check-ups, mental health support, and interventions to manage chronic conditions can mitigate the factors contributing to malnutrition.
4. **Physical Activity Programs:** Encouraging physical activity can help maintain muscle mass and stimulate appetite, reducing the risk of inflammation-induced tissue catabolism.

Conclusion

Among elderly adults who live alone, malnutrition is still a severe problem because of anorexia-induced inflammation-induced tissue catabolism and nutrient deficiency. A comprehensive strategy, including dietary education, community assistance, medical and psychiatric care, and physical activity promotion, is needed to address this complicated issue. By implementing these tactics, we can lessen the toll that malnutrition takes on people and healthcare systems while improving senior citizens' health and quality of life.

References

1. Agarwal E, Miller M, Yaxley A, Isenring E. Malnutrition in older people: a narrative review. *Maturitas*. 2013 Dec;76(4):296-302. doi: 10.1016/j.maturitas.2013.07.013. Epub 2013 Aug 2. PMID: 23958435.
2. Cederholm T, Bosaeus I, Barazzoni R, Bauer J, Van Gossum A, Klek S, Muscaritoli M, Nyulasi I, Ockenga J, Schneider SM, de van der Schueren MA, Singer P. Diagnostic criteria for

- malnutrition - An ESPEN Consensus Statement. *Clin Nutr.* 2015 Jun;34(3):335-40. doi: 10.1016/j.clnu.2015.03.001. Epub 2015 Mar 9. PMID: 25799486.
3. American College of Sports Medicine; Chodzko-Zajko WJ, Proctor DN, Fiatarone Singh MA, Minson CT, Nigg CR, Salem GJ, Skinner JS. American College of Sports Medicine position stand. Exercise and physical activity for older adults. *Med Sci Sports Exerc.* 2009 Jul;41(7):1510-30. doi: 10.1249/MSS.0b013e3181a0c95c. PMID: 19516148.
 4. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, Martin FC, Michel JP, Rolland Y, Schneider SM, Topinková E, Vandewoude M, Zamboni M; European Working Group on Sarcopenia in Older People. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing.* 2010 Jul;39(4):412-23. doi: 10.1093/ageing/afq034. Epub 2010 Apr 13. PMID: 20392703; PMCID: PMC2886201.
 5. Hannan MT, Felson DT, Dawson-Hughes B, Tucker KL, Cupples LA, Wilson PW, Kiel DP. Risk factors for longitudinal bone loss in elderly men and women: the Framingham Osteoporosis Study. *J Bone Miner Res.* 2000 Apr;15(4):710-20. doi: 10.1359/jbmr.2000.15.4.710. PMID: 10780863.
 6. Franceschi C, Bonafè M, Valensin S, Olivieri F, De Luca M, Ottaviani E, De Benedictis G. Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci.* 2000 Jun;908:244-54. doi: 10.1111/j.1749-6632.2000.tb06651.x. PMID: 10911963.
 7. Leslie, W., & Hankey, C. (2015). Aging, Nutritional Status and Health. *Healthcare (Basel, Switzerland)*, 3(3), 648–658. <https://doi.org/10.3390/healthcare3030648>
 8. Locher, J. L., Ritchie, C. S., Roth, D. L., Baker, P. S., Bodner, E. V., & Allman, R. M. (2005). Social isolation, support, and capital and nutritional risk in an older sample: ethnic and gender differences. *Social science & medicine* (1982), 60(4), 747–761. <https://doi.org/10.1016/j.socscimed.2004.06.023>